


FILL PK PLUS

IMPROVES FRUIT QUALITY MODERATES VEGETATIVE LUSHNESS PREPARES PLANTS FOR WINTER REST

With its specific PK 1:1,3 ratio, FILL PK PLUS is formulated to favor fruit and wood ripening, to contain the vegetation development and to prevent undesired early sprouting. Its application during the final phases of the crop cycle prepares the plant to produce and accumulate dry matter, in order for fruit bearing plants to be ready for winter rest.

CROP	TIME OF APPLICATION	DOSE/HECTARE*
Kiwifruit e Grapes	From pre-veraison (change of color) to ripening	4-5 kg
Citrus (Orange, Bergamot, Clementine, Lemon, Tangerine) e Olive	From pre-veraison (change of color) to ripening	4-5 kg
Stone fruits (Apricot, Cherry, Nectarine, Peach, Plum) e Pome fruits (Quince, Apple, Pear)	From pre-veraison (change of color) to ripening	4-5 kg
Strawberries e Small fruits (Raspberry, Blueberry, Blackberry, Currant)	At fruit development	4-5 kg
Fruiting vegetables (Watermelon, Cucumber, Eggplant, Melon, Pepper, Tomato, Zucchini, Pumpkin)	From fruit development to pre-veraison (change of color)	4-5 kg
Industrial crops (Beets, Sugarcane, Rapeseed, Cotton, Sunflower, Industrial tomato, Soybeans, Tobacco)	From fruit development to pre-veraison (change of color)	4-5 kg

COMPOSITION	
Phosphoric anhydride (P ₂ O ₅) soluble in water	40.00%
Phosphoric anhydride (P ₂ O ₅) soluble in neutral ammonium citrate and in water	40.00%
Potassium oxide (K ₂ O) soluble in water	52.00%

PHYSICO-CHEMICAL FEATURES	
SOLUBLE POWDER	
pH (sol 1%)	9.00
Conductivity E.C. S/cm (1‰)	1200
METHOD OF USE	
	Foliar fertilization

PACKAGING: 5 - 25 KG